

# Isodine<sup>®</sup> Medicated Foam Hand Wash F

## Precautions

### Never do

(If you do not follow these instructions, the current symptoms may worsen or adverse reactions are more likely to occur.)

Do not use the following persons.

- (1) Persons who have had allergic symptoms due to this product or any of the ingredients of the product.
- (2) The affected area is widespread.
- (3) Persons have a deep wound or severe burn.

### Consultation

#### 1. Consult a physician, a pharmacist, or a registered sales clerk before using this product if

- (1) Persons are receiving medical treatment.
- (2) You or any of your family members is allergic.
- (3) Persons have had allergic symptoms caused by medicines.

#### 2. In the following cases, immediately stop using this product and consult a physician or a pharmacist, bringing this package insert.

- (1) If any of the following symptoms appears after using this product

Affected site	Symptoms
Skin	Rash/redness, itching

The following serious symptoms may rarely appear.

In such case, immediately consult a physician.

Name of Symptom	Symptoms
Shock (Anaphylaxis)	Immediately after using this product, pale face, cold hands and feet, cold sweat or respiratory distress as well as symptoms such as hives, edema and tightness in the chest may occur.

- (2) If symptoms do not improve despite use of this product for 5 to 6 days.

## Indications

Sterilization/disinfection of hands/fingers and skin

## Dosage and Administration

Take an appropriate amount of this product on your hand, and apply or rub.

### <Precautions related to Dosage and Administration>

#### ● Rinse with water.

- Use this product exactly as indicated under "Dosage and Administration".
- Children should be supervised by their guardians while using this product.
- Take care so that this product does not get into the eyes. Should it get into the eyes, immediately rinse them with water or lukewarm water and see an ophthalmologist if you develop a severe irritation.
- **This product is only for external use; do not ingest or use for gargling.**
- Since the sterilization/disinfection effect of this product may be reduced when it is used in combination with soap etc., wash off soap content before using this product.

## Composition

Each mL of this product contains 75 mg of povidone iodine (available iodine 7.5 mg).

As additives, this product contains sodium lauryl sulfate, dioctyl sodium sulfosuccinate and pH adjuster.

## Precautions for storage and handling

- (1) Store in a cool place avoiding direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer to a different container. (It may lead to misuse or alter the quality.)
- (4) If this product comes into contact with clothes etc., immediately wash it off with water.
- (5) **Since this product discolors products containing silver, remove precious metals before using this product.**
- (6) Do not use the product after the expiration date.

- Disclaimer regarding provision of product information on OTC drugs in foreign languages
- This product is a drug (or designated quasi-drug) approved based on the Act on Securing Quality, Efficacy and Safety of Products Including Pharmaceuticals and Medical Devices in Japan for sale and use in Japan.
- The product information in foreign languages is a translation of the product's package insert written in Japanese, and provided just as supporting information. It is not guaranteed that the information contained and the product itself comply with laws and regulations outside Japan.
- The product information in foreign languages is a tentative translation by the company, and no warranty is given to the translated content. It is subject to modification/change without prior notice in the future.
- We assume no responsibility whatsoever for any problems attributable to the content of the product information in foreign languages.